Recent results on the measurements of the P-odd correlations in the capture of slow polarized neutrons by  $^6\mathrm{Li}$  and  $^{10}\mathrm{B}$ 

Yuri M. Gledenov<sup>1</sup>, Victor A. Vesna<sup>2</sup>, Ekaterina V. Shulgina<sup>2</sup>, Pavel V. Sedyshev<sup>1</sup>, Valeriy V. Nesvizhevsky<sup>3</sup>, Alexander K. Petukhov<sup>3</sup>, Torsten Soldner<sup>3</sup>, Oliver Zimmer<sup>4</sup>

Two experiments on the P-odd correlation measurement were carried out at the high intensity polarized cold neutron beam of the PF1B instrument of the ILL reactor, Grenoble, France. The triton emission asymmetry in the  $^6Li(n,\alpha)^3H$  reaction and the asymmetry of the emitted  $\gamma$ -quanta at discharge of the  $^7Li$  first excited state populated in the  $^{10}B(n,\alpha)^7Li$  reaction were measured. In the experiments a special apparatus and measurement method were used. The preliminary results of the  $^6Li$  and  $^{10}B$  measurements are  $\alpha^t_{PNC} = -(8.1 \pm 3.9)10^{-8}$  and  $\alpha^{\gamma}_{PNC} = -(11.0 \pm 6.6)10^{-8}$ , respectively. At the base of first result, the most accurate at present day estimation of the weak neutral current constant has been given:  $f_{\pi} \leq 1.210^{-7}$ . The results are discussed.

Email: gledenov@nf.jinr.ru

<sup>&</sup>lt;sup>1</sup> Joint Institute for Nuclear Research

<sup>&</sup>lt;sup>2</sup> Petersburg Nuclear Physics Institute

 $<sup>^3</sup>$  Institute Laue-Langevin

<sup>&</sup>lt;sup>4</sup> Technishe Uneversitat Munchen